

1. THE DIGESTIVE SYSTEM

NOW PLEASE COMPLETE THE FOLLOWING SUBQUESTIONNAIRES USING THE SAME RATING SYSTEM:
Leave blank if symptom or activity does not apply, 1 for mild or rarely occurring, 2 for moderate or regularly occurring,
3 for severe or often occurring.

UNDERACTIVE STOMACH

Excessive gas, belching or burping after meals	
Stomach bloated after eating	
Sleepy after eating	
Longitudinal striations on fingernails	
Eat when rushed/in a hurry	
Halitosis	
Full feeling after heavy meat meal	
Heavy, tired feeling after eating	
Nausea after taking supplements	
Acne	
Undigested food in the stool	

GALL BLADDER:

Gall stones; history of gall stones	
Stool appears clay-coloured, foul odoured	
Constipation	
High cholesterol diet; High blood cholesterol levels	
Severe pain in right upper abdomen	

LIVER

Yellow or pale fingernails	
Skin oily on nose and forehead	
Fats/greasy foods cause nausea, headaches	
Vertical white streaks on fingernails	
Onions, cabbage, radishes, cucumbers cause bloating /gas	
Bad breath; bad taste in mouth	
Excess body odour	
High cholesterol / high cholesterol diet	
Stiff, aching muscles	
Migraine headaches	
Discomfort underneath right ribcage	
Food allergies	
Irritable, easily angered	
Weight gain around the abdomen	
Yellow palms	
Jaundice	
Poor concentration	
Difficulty losing weight	
Acne, boils, rashes, psoriasis or eczema	
Constipation	

OVERACTIVE STOMACH

Stomach pain 1 hour after eating or at night	
Burning sensation in stomach	
Pain aggravated by worry / tension	
Hiatal hernia	
Gastritis, gastric ulcer	
Nausea, vomiting	
Sensation of acidity in abdominal area	
Heartburn, indigestion	
Blood in stool	
Lower back pain	
Long term aspirin use	

PANCREAS

Severe abdominal pain	
Nausea and vomiting	
Slow digestion; feel full for hours after eating	
Fever	
Alcohol addiction	
Jaundice	

DYSGLYCEMIA

Hungry up to 3 hours after eating	
Strong, sudden cravings for sweets, starches coffee or alcohol	
Nervous/anxious feelings relieved by eating	
Irritable if late for, or skip, a meal	
Overweight	
Addicted to coffee with sugar and/or colas	
Frequent "midnight snacks"	
Family history of diabetes	
Fatigue	
Frequent headaches	
Fainting spells	
Depression	
Lose temper easily	